

REFERENCES

- ACCEO. 2003. The Alliance for California Current Observation. Available at <http://swfsc.nmfs.noaa.gov/FRD/acceo/acceo1.htm>. (Accessed November 2004].
- Barbour, M. G., and J. Major (eds.). 1977. Terrestrial vegetation of California. Wiley, New York, 1002 p.
- Bisson, P. A., K. Sullivan, and J. L. Nielsen. 1988. Channel hydraulics, habitat use, and body form of juvenile coho salmon, steelhead and cutthroat trout in streams. *Trans. Am. Fish. Soc.* 117:262-273.
- Bjorkstedt, E. 2004. Ranking putative populations by relative abundance and isolation: a tool for guiding delineation of historical independent populations based on simple demographic considerations. NOAA/NMFS/SWFSC. 11 p.
- Brodeur, R. D., R. L. Emmett, J. P. Fisher, E. Casillas, D. J. Teel, and T. W. Miller. 2004. Juvenile salmonid distribution, growth, condition, origin, and environmental and species associations in the Northern California Current. *Fish. Bull.* 102:25-46.
- Brodeur, R. D., R. L. Emmett, J. P. Fisher, E. Casillas, D. J. Teel, and T. W. Miller. 2003. Juvenile salmonid distribution, growth, condition, origin, and environmental and species associations in the Northern California Current. *Fish. Bull.* 102:25-46.
- Buckling, A., M. A. Wills, and N. Colgrave. 2003. Adaptation limits diversification of experimental bacteria populations. *Science* 302:2107-2109.
- Buffington, J. M., T. E. Lisle, R. D. Woodsmith, and S. Hilton. 2002. Controls on the size and occurrence of pools in coarse-grained forest rivers. *River Res. Appl.* 18:507-531.
- Burnett, K. 2001. Relationships among juvenile anadromous salmonids, their freshwater habitat, and landscape characteristics over multiple years and spatial scales in the Elk River, Oregon. Ph.D., Oregon State University. Available at <http://www.fsl.orst.edu/clams/pubs.html>.
- Burnett, K. September 2003. Personal Communication. Fish Biologist, USDA Forest Service, Pacific Northwest Research Station, Aquatic and Land Interaction Program. (Available from K. Burnett, 3200 Jefferson Way, Corvallis, OR 97331.)
- Burnett, K. M., G. H. Reeves, D. Miller, S. E. Clarke, K. C. Christiansen, and K. Vance-Borland. 2003. A first step toward broad scale identification of freshwater protected areas for Pacific salmon and trout, p. 144-154. *In* J. Beumer, A. Grant, and D. C. Smith (eds.), *Aquatic Protected Areas: what works best and how do we know?*, Proceedings of the World Congress on Aquatic Protected Areas, Cairns, Australia, August 14-18, 2002. Australian Society for Fish Biology, University of Queensland, Australia.
- Cedarholm, J. C., M. D. Kuntz, T. Murota, and A. Sibatani. 1999. Pacific salmon carcasses: essential contributions of nutrients and energy for aquatic and terrestrial systems. *Fisheries* 24:6-13.
- Chapman, D. W. 1965. Net production of juvenile coho salmon in three Oregon streams. *Trans. Am. Fish. Soc.* 94(1):40-52.

- Chapman, D. W. 1986. Salmon and steelhead abundance in the Columbia River in the nineteenth century. *Trans. Am. Fish. Soc.* 115:662-670.
- Coulter, K., P. Williams, and P. Benner. 1996. An Environmental History of the Tillamook Bay National Estuary Project Technical Report 09-96. USEPA #CE 99020292-1.
- Currens, K. P., and D. Farnsworth. 1993. Mitochondrial DNA variation in Oregon coho salmon. Report to Oregon Department of Fish and Wildlife, 16 p. (Available from Oregon Department of Fish and Wildlife, 3406 Cherry Ave. NE, Salem, OR 97303.)
- Dennis, B., P. Mulholland, and J. M. Scott. Estimation of Growth and Extinction Parameters for Endangered Species. *Ecol. Monogr.* 61(2):115-143.
- Dose, J. 2004. Personal Communication. Umpqua National Forest, 2900 Stewart Parkway, Roseburg, OR 97470. 541/ 672-6601. Jan. 30, 2004.
- Drucker, B. 1972. Some life history characteristics of coho salmon of the Karluk River system, Kodiak Island, Alaska. *Fish. Bull.* 70:79-94.
- Eames, M., T. Quinn, K. Reidinger, and D. Haring. 1981. Northern Puget Sound 1976 adult coho and chum tagging studies. *Wash. Dep. Fish. Tech. Rep.* 64, 136 p.
- Ehrlich, P., and J. Roughgarden. 1987. *The Science of Ecology.* McMillan, New York. 710 p.
- Flagg, T. A., F. W. Waknitz, D. J. Maynard, G. B. Milner, and C. V. Mahnken. 1995. The effect of hatcheries on native coho salmon populations in the lower Columbia River. In H. Schramm and B. Piper (eds.), *Proceedings of the American Fisheries Society Symposium on the uses and effects of cultured fishes in aquatic ecosystems*, March 12-17, 1994, Albuquerque, NM. *Am. Fish. Soc. Symp.* 15:366-375.
- Flint, T., and G. Zillges. 1980. Little Bear Creek coho salmon stream life study. *Wash. Dep. Fish. Prog. Rep.* 124, 40 p.
- Forbes, S., K. Knudsen, and F. Allendorf. 1993. Genetic variation in DNA of coho salmon from the Lower Columbia River. Final Report to the U.S. Dep. Energy, Bonneville Power Administration, Division of Fish and Wildlife, Contract DE-BI79-92BP 30198, 25 p. (Available from U.S. Department of Energy, Bonneville Power Administration, Public Information Office - ALP-22, P.O. Box 3621, Portland, OR 97208.)
- Ford, M. J., D. Teel, D. M. van Doornik, D. Kuligowski, and P. W. Lawson. 2004. Genetic population structure of Central Oregon Coastal coho salmon (*Oncorhynchus kisutch*). Unpublished draft report. 35 p. (Available from NOAA Fisheries, Northwest Fisheries Science Center, 2725 Montlake Blvd. East, Seattle WA 98112-2097.)
- Franklin, J. F., and C. T. Dyrness. 1973. Natural vegetation of Oregon and Washington. USDA Forest Service General Tech. Rep. PNW-8, 417 p. U.S. Gov. Printing Office, Washington, DC.
- Gilbert, C. H. 1912. Age at maturity of the Pacific coast salmon of the genus *Oncorhynchus*. *Fish Bull.* 32:3-22.
- Godfrey, H. 1965. Coho salmon. *Int. North Pac. Fish. Comm. Bull.* 16, 39 p.

- Good, J. 2000. Summary and current status of Oregon's estuarine ecosystems. *In* State of the Environment Report. (Available from Oregon Progress Board, 155 Cottage Street NE, Salem, OR 97301.)
- Government of Canada. 2003a. The Atlas of Canada. National Frameworks Administrative Boundaries, Canada. [Vector digital data online]. Available at <http://geogratis.gc.ca/download/frameworkdata/admin/Canada/bnd.e00.gz>. (Accessed 5 November, 2003.)
- Government of Canada. 2003b. Natural Resources Canada. Canadian Geographical Names – Querying Canadian Geographical Names. [Online]. Available at http://geonames.nrcan.gc.ca/search/search_e.php. (Accessed 5 November, 2003.)
- Gregory, S. V., G. A. Lamberti, and K. M. S. Moore. 1989. Influence of valley floor land forms on stream ecosystems, p. 3-8. *In* D. L. Abell (ed.), Proceedings of the California riparian systems conference: Protection, management, and restoration for the 1990s, 22-24 September 1988, Davis, California. USDA Forest Service General Tech. Rep. PSW-110. Pacific Southwest Forest and Range Experiment Station, Berkeley, CA.
- Hanski, I. A., and M. E. Gilpin (eds.). 1997. Metapopulation Biology: ecology, genetics, and evolution. Academic Press, San Diego. 512 p.
- Hartman, G. F., L. B. Holtby, and J. C. Scrivener. 1984. Some effects of natural and logging-related winter stream temperature changes on the early life history of coho salmon (*Oncorhynchus kisutch*) in Carnation Creek, British Columbia. *In* W. R. Meehan, T. R. Merrill, Jr., and T. A. Hanley (eds.), Proceedings of a symposium, Fish and wildlife relationships in old-growth forests, 12-15 April 1982, Juneau, Alaska, p. 141-149. Alaska District, American Institute of Fishery Research Biologists, Northwest Section, The Wildlife Society, Alaska Council on Science and Technology,
- Hartt, A. C. 1980. Juvenile salmonids in the oceanic ecosystem – the critical first summer. *In* W. J. McNeil and D. C. Himsworth (eds.), Salmonid ecosystems of the North Pacific, p. 25-57. Oreg. State Univ. Press and Oreg. State Univ. Sea Grant College Program, Corvallis, OR.
- Hartt, A. C., and M. B. Dell. 1986. Early oceanic migrations and growth of juvenile Pacific salmon and steelhead trout. *Int. North Pac. Fish. Comm. Bull.* 46, 105 p.
- Hickman, J. C. 1993. The Jepson manual: higher plants of California. University of California Press, Berkeley, CA. 1400 p.
- Hilborn, R., and C. J. Walters. 1992. Quantitative Fisheries Stock Assessment: Choice, Dynamics & Uncertainty. Chapman and Hall, New York. 570 p.
- Hjort, R. C., and C. B. Schreck. 1982. Phenotypic differences among stocks of hatchery and wild coho salmon, *Oncorhynchus kisutch*, in Oregon, Washington, and California. *Fish. Bull.* 80:105-11.
- Holtby, L. B. 1987. The effects of logging on the coho salmon of Carnation Creek, British Columbia. *In* T. W. Chamberlin (ed.), Proceedings of the workshop: Applying 15 years of Carnation Creek results, January 13-15, 1987, Pacific Biological Station, Nanaimo, British Columbia, p. 159-174. Carnation Creek Steering Committee, British Columbia. (Available from J. C. Scrivener, Pacific Biological Station, Box 100, Nanaimo, B. C., Canada V9R 5K6.)

- Huntington, C. 2003. Preliminary draft Oregon Coast coho populations. (Unpublished data). Compiled from Kostow 1995.
- Hydrosphere Data Products, Inc. 1995. Hydrodata Regional CD-ROMs: U.S. Geological Survey daily values, Vols. West 1, West 2. In L. A. Weitkamp, et al. (eds.), Status Review of Coho Salmon from Washington, Oregon, and California. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-NWFSC-24, 258 p.
- Iwamoto, R. N., B. A. Alexander, and W. K. Hershberger. 1984. Genotypic and environmental effects on the incidence of sexual precocity in coho salmon (*Oncorhynchus kisutch*). Aquaculture 43(1984):105-121.
- IMST (Independent Multidisciplinary Science Team). 2002. Recovery of Wild Salmonids in Western Oregon Lowlands. A report of the Independent Multidisciplinary Science Team, Oregon Plan for Salmon and Watersheds. Tech. Rep. 2002-1 to the Oregon Plan for Salmon and Watersheds, Governor's Natural Resources Office, Salem, Oregon. 169 p.
- Johnson, O. W., T. A. Flagg, D. J. Maynard, G. B. Milner, and F. W. Waknitz. 1991. Status review for lower Columbia River coho salmon. U.S. Dep. Commer., NOAA Tech. Memo. NMFS F/NWC-202, 94 p.
- Johnson, S. L. 1988. The effects of the 1983 El Niño on Oregon's coho (*Oncorhynchus kisutch*) and chinook (*O. tshawytscha*) salmon. Fish. Res. 6:105-123.
- Johnson, S. L., M. F. Solazzi, and J. D. Rogers. 1993. Development and evaluation of techniques to rehabilitate Oregon's wild salmonids. Oreg. Dep. Fish Wildl. Fish Research Proj. F-125-R-5, Annual Rep, 12 p. (Available from Oregon Department of Fish and Wildlife, P.O. Box 59, Portland, OR 97207.)
- Knight, N. J. 1979. Factors affecting the smolt yield of coho salmon (*Oncorhynchus kisutch*) in three Oregon streams. M.S. Thesis, Oregon State Univ., 105 p.
- Kostow, K. 1995. Biennial Report on the Status of Wild Fish in Oregon. Oregon Department of Fish and Wildlife, 213 p. (Report Available from ODFW 3406 Cherry Ave. NE, Salem, OR 97303).
- KCM (Kramer, Chin, and Mayo, Inc.). 1983. Devils Lake diagnostic and feasibility study. (Available from Lincoln City Planning Dept., 801 SW Hwy 101, Lincoln City, OR 97367.)
- Lande, R. 1993. Risks of population extinction from demographic and environmental stochasticity and random catastrophes. Am. Nat. 142(6):911-927.
- Lande, R. 1998. Demographic stochasticity and Allee effect on a scale with isotropic noise. Oikos 83:353-358.
- Lawson, P. W., E. A. Logerwell, N. J. Mantua, R. C. Francis, and V. N. Agostini. 2004. Environmental factors influencing freshwater survival and smolt production in two Pacific Northwest coho salmon (*Oncorhynchus kisutch*) populations. Can. J. Fish. Aquat. Sci. 61:360-373.
- Lichatowich, J. A. 1989. Habitat alteration and changes in abundance of coho (*Oncorhynchus kisutch*) and chinook salmon (*O. tshawytscha*) in Oregon's coastal streams. In C. D. Levings, L. B.

- Holtby, and M. A. Henderson (eds.) ,Proceedings of the national workshop on effects of habitat alteration on salmonid stocks. Can. Spec. Publ. Fish. Aquat. Sci. 105, p. 92-99.
- Lister, D. B., L. M. Thorson, and I. Wallace. 1981. Chinook and coho salmon escapements and coded-wire tag returns to the Cowichan-Koksilah river system, 1976-1979. Can. Manuscript Rep. Fish. Aquat. Sci. 1608, 168 p.
- Loeffel, R. E., and H. O. Wendler. 1968. Review of the Pacific coast chinook and coho salmon resources with special emphasis on the troll fishery. Prepared by the U.S. working group of the Informal Committee on Chinook and Coho, 107 p. (Available from Oregon Department of Fish and Wildlife, P.O. Box 59, Portland, OR 97207.)
- Loeffel, R. E., and W. O. Forster. 1970. Determination of movement and identity of stocks of coho salmon in the ocean using the radionuclide Zinc-65. Res. Rep. Fish Comm. Oregon, 2(1):1-13.
- Mangel, M., and C. Tier. 1994. Four facts every conservation biologist should know about persistence. Ecology 75 (3):607-614.
- Mayr, E. 1970. Populations, Species, and Evolution. Harvard University Press, Cambridge, Mass. 453 p.
- McElhany, P., M. H. Ruckelshaus, M. J. Ford, T. C. Wainwright, and E. P. Bjorkstedt. 2000. Viable salmonid populations and the recovery of evolutionarily significant units. U.S. Dept. Commer., NOAA Tech. Memo. NMFS-NWFSC-42, 156 p.
- McGie, A. M. 1970. Research and management studies of Oregon coastal salmon. Annual Report, July 1, 1969, to June 30, 1970. Fish. Comm. Oregon, Div. Manage. Res., Coastal Rivers Invest., 107 p. (Available from Oregon Department of Fish and Wildlife, 3406 Cherry Ave. NE., Salem, OR 97303.)
- McGie, A. M. 1971. Salmon research on coastal streams and reservoirs. Annual Report July 1, 1970 to June 30, 1971. Fish. Comm. Oregon, Div. Manage. Res., Coastal Rivers Invest., 55 p. (Available from Oregon Department of Fish and Wildlife, 3406 Cherry Ave. NE., Salem, OR 97303.)
- McGie, A. M. 1972. Salmon research on coastal streams and reservoirs. Annual Report, July 1, 1971 to June 30, 1972. Fish. Comm. Oregon, Div. Manage. Res., Coastal Rivers Invest., 41 p. (Available from Oregon Department of Fish and Wildlife, 3406 Cherry Ave. NE., Salem, OR 97303.)
- McGie, A. M. 1973. Salmon research on coastal streams and reservoirs. Annual Report July 1, 1972 to June 30, 1973. Fish. Comm. Oregon, Div. Manage. Res., Coastal Rivers Invest., 31 p. (Available from Oregon Department of Fish and Wildlife, 3406 Cherry Ave. NE., Salem, OR 97303.)
- McGie, A. M. 1975. Research on anadromous fish in coastal watersheds of Oregon. Annual Report, July 1, 1974 to June 30, 1975. Fish. Comm. Oregon, Div. Manage. Res., Coastal Rivers Invest., 19 p. (Available from Oregon Department of Fish and Wildlife, 3406 Cherry Ave. NE., Salem, OR 97303.)

- McGie, A. M. 1977. Anadromous fish research in Oregon's coastal watersheds. Oreg. Fish Commission Federal Aid Progress Rep., Fisheries. Fish. Comm. Oregon, Div. Manage. Res., Coastal Rivers Invest., 55 p. (Available from Oregon Department of Fish and Wildlife, 3406 Cherry Ave. NE, Salem, OR 97303.)
- McNaughton, S. J., and L. L. Wolf. 1973. General Ecology. Holt, Reinhardt and Winston, Inc., New York. 710 p.
- Meffe, G. K., and R. C. Vrijenhoek. 1988. Conservation genetics in the management of desert fishes. *Conserv. Biol.* 2:157-169.
- Miller, D. 2002. Programs for DEM analysis. Available at <http://www.orst.edu/clams>.
- Miller, D. R., J. G. Williams, and C. W. Sims. 1983. Distribution, abundance and growth of juvenile salmonids off the coast of Oregon and Washington, summer 1980. *Fish. Res.* 2:1-17.
- Milner, G. B. 1993. Isozyme variation of coho salmon (*Oncorhynchus kisutch*) and its potential to estimate stock compositions of mixed stock fisheries. In L. Berg and P. W. Delaney (eds.), *Proceedings of the Coho Workshop*, Nanaimo, B. C., May 26-28, 1992, p. 182-192. (Available from Canadian Department of Fisheries and Oceans, Habitat Management Sector, Policy and Information Unit, 327-555 W. Hastings St., Vancouver, B.C. V6B 5G3.)
- Montgomery, D. R., and J. M. Buffington. 1997. Channel-reach morphology in mountain drainage basins. *Geol. Soc. Am. Bull.* 109:596-611.
- Morgan, A. R., and K. A. Henry. 1959. The 1955-1956 silver salmon run into the Tenmile Lakes System. *Fish Comm. Oreg., Res. Briefs* 7(1):57-77.
- Moring, J. R., and R. L. Lantz. 1975. The Alsea watershed study: effects of logging on the aquatic resources of three headwater streams of the Alsea River, Oregon. Part I- Biological studies. *Oreg. Dep. Fish Wildl., Fish. Res. Rep.* 9, 66 p.
- Moritz, C., S. Lavery, and R. Slade. 1995. Using allele frequency and physiology to define units for conservation and management. In J. L. Nielsen (ed.), *Symposium on Evolution and the Aquatic Ecosystem: Defining Unique Units in Population Conservation*, Monterey, CA (USA), 23-25 May 1994. *Am. Fish. Soc. Symp.* 17:249-262.
- Mullen, R. E. 1981a. Oregon's commercial harvest of coho salmon *Oncorhynchus kisutch* (Walbaum). *Oreg. Dep. Fish and Wildl., Fish Div. Info. Rep.* 81-3. (App. III.)
- Mullen, R. E. 1981b. Estimates of historical abundance of coho salmon *Oncorhynchus kisutch* (Walbaum), in Oregon coastal streams and in the Oregon Production Index area. *Oreg. Dep. Fish and Wildl., Fish Div. Info. Rep.* 81-5. (App. III.)
- Myers, J., C. Busak, D. Rawding, and A. Marshal. 2003. Historical population structure of Willamette and Lower Columbia River Basin Pacific salmonids. Willamette-Lower Columbia-TRT Report. NOAA Fisheries Northwest Fisheries Science Center, Seattle, Washington.
- Naiman R. J., T. J. Beechie, L. E. Benda, D. R. Berg, P. A. Bisson, L. H. McDonald, M. D. O'Connor, P. L. Olson, and E. A. Steel. 1992. Fundamental elements of ecologically healthy watersheds in

- the Pacific Northwest Coastal Ecoregion. In R. J. Naiman (ed.), *Watershed Management: Balancing Sustainability and Environmental Change*, p. 127-188. Springer-Verlag, New York.
- Nicholas, J. W., L. Van Dyke, and R. C. Buckman. 1982. Straying by hatchery-reared coho salmon released in Yaquina Bay, Oregon. *Oreg. Dep. Fish Wildl., Info. Rep.* 82-6, 22 p.
- Nickelson, T. E. 2001. Population Assessment: Oregon Coast Coho Salmon ESU. Oregon Dept. Fish Wildl. Info. Rep. 2001-02. 47 p.
- Nickelson, T. E. 1998. A habitat-based assessment of coho salmon production potential and spawner escapement needs for Oregon coastal streams. *Oreg. Dep. Fish Wildl., Fish Div. Info. Rep.* 98-4. (App. III.)
- Nickelson, T. E., and P. W. Lawson. 1998. Population viability of coho salmon (*Oncorhynchus kisutch*) in Oregon coastal basins: application of a habitat –based life history model. *Can. J. Fish. Aquat. Sci.* 55:2383-2392.
- Nielsen, J. L. 1994. Invasive cohorts: Impacts of hatchery-reared coho salmon on the trophic, developmental, and genetic ecology of wild stocks. In D. J. Stouder, K. L. Fresh, R. J. Feller (eds.), *Theory and application in fish feeding ecology*, Vol. 18 of The Belle W. Baruch Library in Marine Science, p. 361-385. Univ. S. Carolina Press, Columbia.
- NMFS (National Marine Fisheries Service). 1991. Notice of Policy: Policy on applying the definition of species under the Endangered Species Act to Pacific salmon. *Fed. Reg.* 56:58612-58616.
- NMFS (National Marine Fisheries Service), Quinalt Treaty Area Indian Tribes, U.S. Fish and Wildlife Service (USFWS), and Washington State Department of Fisheries (WDF). 1983. Coho salmon spawning escapement goals for North Washington coastal rivers. Final Report of a workshop held October 13 & 14 1982 in Renton, WA. (cited in Appendix III)
- NOAA Fisheries (National Oceanic and Atmospheric Administration Fisheries). 1999. (No title given) [Vector digital data online]. Available at http://www.nwr.noaa.gov/iesumaps/cohoesu_e00.zip. (Accessed 2 July 2003.)
- NOAA Fisheries. 2003. Transmittal letter from NWFSC to Regional Office.
- NOAA Fisheries. 2004. Oregon Coast coho run timing for 2001. ODFW unpublished data. Available at <http://www.nwfsc.noaa.gov/trt/orcoast.htm#docs>. (Accessed 30 January 2004.)
- ODFW (Oregon Department of Fish and Wildlife). 1995. Biennial Report on the Status of Wild Fish in Oregon. Oregon Department of Fish and Wildlife, Portland, OR. 217 p.
- ODFW (Oregon Department of Fish and Wildlife). 2002a. Natural Resources Information Management Program. Barriers to Fish Passage. [Vector digital data online]. Available at ftp://rainbow.dfw.state.or.us/pub/gis/other/shape/bar_pts.zip. (Accessed 29 August 2003.)
- ODFW (Oregon Department of Fish and Wildlife). 2002b. Natural Resources Information Management Program (2002). ODFW Dams Coverage. Available at ftp://rainbow.dfw.state.or.us/pub/gis/other/shape/dam_pts.zip. (Accessed 30 January 2004.)

- ODFW (Oregon Department of Fish and Wildlife). 2003a. Natural Resources Information Management Program. Distribution of Oregon anadromous salmon and steelhead habitat: coho salmon. Eleventh edition. [Vector digital data online]. Available at ftp://rainbow.dfw.state.or.us/pub/gis/k100/shape/coho/coho_v11.zip. (Accessed 16 July 2003.)
- ODFW (Oregon Department of Fish and Wildlife). 2003b. Homing Fidelity Calculations. Unpublished Data. T. Nickelson, Personal Communication. Corvallis Research Laboratory, 28655 Hwy 34, Corvallis, OR 97333.
- ODFW (Oregon Department of Fish and Wildlife). 2003c. Potential Historical Abundance. ODFW Research Website. Available at [http://oregonstate.edu/dept/odfw/coastal_salmonid inventories/coho abundance/estimates for coho lake basins](http://oregonstate.edu/dept/odfw/coastal_salmonid_inventories/coho_abundance/estimates_for_coho_lake_basins). (Accessed 10 September 2003.)
- ODFW (Oregon Department of Fish and Wildlife). 2003d. Lake System tributary spawning run for 2001. Available at [http://Oregonstate.edu/Dept/ODFW/Coastal salmonid inventories/coho abundance/estimates for coho lake basins](http://Oregonstate.edu/Dept/ODFW/Coastal_salmonid_inventories/coho_abundance/estimates_for_coho_lake_basins). (Accessed 15 September 2003.)
- Olin, P. G. 1984. Genetic variability in hatchery and wild populations of coho salmon, *Oncorhynchus kisutch*, in Oregon. M.S. Thesis, Univ. California Davis. 73 p.
- ONHP (Oregon Natural Heritage Program). 2001. ORNHP / ODFW Ecoregions of Oregon. [Vector digital data online]. Oregon Geospatial Data Clearinghouse. Available at <http://www.sscgis.state.or.us/data/shapefile/k250/ecoregion.zip>. (Accessed July 31, 2003.)
- ONHP (Oregon Natural Heritage Program). 2003. Oregon Natural Heritage Plan. (Available from Oregon Department of State Lands, 775 Summer Street NE, Suite 100 Salem, OR 97301-1279.)
- Oregon Climate Service. 2000. United States Average Annual Precipitation, 1961-1990. [Vector digital data online]. NRCS (Natural Resources Conservation Service). Available at ftp://ftp.ftw.nrcs.usda.gov/pub/ams/prism/data/states/or/or_shp.zip. (Accessed 1 August 2003.)
- Orr, E. L., and W. N. Orr. 2000. Geology of Oregon. Kendall/Hunt Publishing Company. Dubuque, Iowa. 254 p.
- Pearcy, W. G. 1992. Ocean ecology of North Pacific salmonids. Univ. Washington Press, Seattle. 179 p.
- Pearcy, W. G., and J. P. Fisher. 1988. Migrations of coho salmon, *Oncorhynchus kisutch*, during their first summer in the ocean. Fish. Bull. 86(2):173-186.
- Pess, G. R., D. R. Montgomery, E. A. Steel, R. E. Bilby, B. E. Feist, and H. M. Greenberg. 2002. Landscape characteristics, land use, and coho salmon (*Oncorhynchus kisutch*) abundance, Snohomish River, Washington, U.S.A. Can. J. Fish. Aquat. Sci. 59:613-623.
- Pritchard, A. L. 1940. Studies on the age of the coho salmon (*Oncorhynchus kisutch*) and the spring salmon (*Oncorhynchus tshawytscha*) in British Columbia. Trans. R. Soc. Can., Ser. 3, 34(V):99-120.
- Proctor, C. M., J. C. Garcia, D. V. Galvin, G. C. Lewis, L. C. Lower, and A. M. Mass. 1980. An ecological characterization of the Pacific Northwest Coastal Region. U.S. Fish and Wildlife Service, Biological Services Program. FWS/OBS-79/11 through 79/15, 5 vol.

- Puget Sound Technical Recovery Team. 2003. Independent populations for chinook salmon in Puget Sound. Draft Report. (Available from National Marine Fisheries Service, Northwest Fisheries Science Center, 2725 Montlake Blvd. E., Seattle, WA 98112.)
- Reeves, G. H. 2003. Personal communication. Research Fish Biologist, USDA Forest Service, Pacific Northwest Research Station, Aquatic and Land Interaction Program, 3200 Jefferson Way, Corvallis, OR 97331.
- Reeves, G. H., L. E. Benda, K. M. Burnett, P. A. Bisson, and J. R. Sedell. 1995. A disturbance-based ecosystem approach to maintaining and restoring freshwater habitats of evolutionarily significant units of anadromous salmonids in the Pacific Northwest. *Am. Fish. Soc. Symp.* 17:334-349.
- Reeves, G. H., P. A. Bisson, and J. M. Debaucher. 1998. Fish communities. *In* R. J. Naiman and R. E. Bilby (eds.), *River ecology and management: lessons from the Pacific Coastal Ecoregion*, p. 200-234. Springer, New York.
- Regional Ecosystem Office. 2002. Hydrologic Unit Boundaries for Oregon, Washington, and California: Version 1.3. [Vector digital data online]. Regional Ecosystem Office. Available at <http://www.reo.gov/gis/projects/watersheds/reohucv13.zip>. (Accessed 11 July 2003.)
- Reimers, P. E. 1971. The movement of yearling coho salmon through Sixes River Estuary. Fish Comm. Oregon, Coastal Rivers Investigation Info. Rep. 71-2, 15 p. (Available from Oregon Department of Fish and Wildlife, 3406 Cherry Ave. NE, Salem, OR 97303.)
- Reimers, P., R. Bender, J. Muck, T. Rumreich, R. Smith, and M. Manion. 1993. Fish management review, Southwest Region, Tenmile, Coos, Coquille District, February 9-10, 1993, p. 1-6. (Available from Oregon Department of Fish and Wildlife, 3406 Cherry Ave. NE, Salem, OR 97303.)
- Ricker, W. E. 1972. Hereditary and environmental factors affecting certain salmonid populations. *In* R. C. Simon and P. A. Larkin (eds.), *The Stock Concept in Pacific Salmon*, p. 19-160. MacMillan Lectures on Fisheries. University of British Columbia, Vancouver, B.C.
- Rieman, B. E., and J. B. Dunham. 2000. Metapopulations and salmonids: a synthesis of life history patterns and empirical observations. *Ecol. Freshwat. Fish.* 9(1-2):51-64.
- Rodgers, J. D., R. D. Ewing, and J. D. Hall. 1987. Physiological changes during seaward migration of wild juvenile coho salmon (*Oncorhynchus kisutch*). *Can. J. Fish. Aquat. Sci.* 44(2):452-457.
- Rodgers, J. D., S. L. Johnson, T. E. Nickelson, and M. F. Solazzi. 1993. The seasonal use of natural and constructed habitat by juvenile coho salmon (*Oncorhynchus kisutch*) and preliminary results from two habitat improvement projects on smolt production in Oregon coastal streams. *In* L. Berg and P. W. Delaney (eds.), *Proceedings of the Coho Workshop*, Nanaimo, B. C., May 26-28, 1992, p. 344-351. (Available from Department of Fisheries and Oceans, Habitat Management Sector, Policy and Information Unit, 327-555 West Hastings Street, Vancouver, BC V6B 5G3.)
- Roper, B. (U.S. Forest Service, Tiller Ranger District). 1994. Letter to L. Weitkamp (NMFS) 5/28/94. Enclosed information reflects current understanding of juvenile coho salmon within the upper south Umpqua Basin, OR.
- Rosgen, D. L. 1994. A classification of natural rivers. *Catena* 22:169-199. (App. III.)

- Rosgen, D. L. 1996. Applied River Morphology. Wildland Hydrology, Pagosa Springs, Colorado. (App.III.)
- Sandercock, F. K. 1991. Life history of coho salmon (*Oncorhynchus kisutch*). In C. Groot and L. Margolis (eds.), Pacific salmon life histories, p. 396-445. Univ. British Columbia Press, Vancouver.
- Sedell, J. R., and W. S. Duvall. 1985. Influence of forest and rangeland management on anadromous fish habitat in Western North America –1. Water transportation and storage of logs. Gen. Tech. Rep. PNW-186. Pacific Northwest Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture, Portland, OR.
- Shaffer, M. L. 1981. Minimum population sizes for species conservation. Bioscience 31:131-134.
- Shapovalov, L., and A. C. Taft. 1954. The life histories of the steelhead rainbow trout (*Salmo gairdneri gairdneri*) and silver salmon (*Oncorhynchus kisutch*) with special reference to Waddell Creek, California, and recommendations regarding their management. Calif. Dep. Fish Game, Fish Bull. 98, 375 p.
- Signell, R. 2003. NGDC/WDC MGG, Boulder-Coastline Extractor. [Vector digital data]. National Oceanic and Atmospheric Administration. Available at <http://rimmer.ngdc.noaa.gov/coast/>. (Accessed 24 November, 2003.)
- Silverstein, J. T., and W. K. Hershberger. 1992. Precocious maturation in coho salmon (*Oncorhynchus kisutch*): Estimation of the additive genetic variance. J. Heredity 83:282-286.
- Skeesick, D. G. 1970. The fall immigration of juvenile coho salmon into a small tributary. Fish Comm. Oregon Res. Rep. 2(1):1-6.
- Solazzi, M. F., S. L. Johnson, B. Miller, T. Dalton, K. A. Leader. 2003. Salmonid life-cycle monitoring project 2002. Monitoring Program Report number OPSW-ODFW-2003-2, Oregon Department of Fish and Wildlife, Portland, OR.
- Solazzi, M. F. 1986. Electrophoretic stock characterization of coho salmon populations in Oregon and Washington, and coastal chinook salmon populations in Oregon. Ore. Dep. Fish Wildl. Info. Rep. 86-5, 16 p.
- Soulé, M. E. (ed.). 1987. Viable populations for conservation. Cambridge University Press, Cambridge. 189 p.
- Steward, C. R., and T. C. Bjornn. 1990. Supplementation of salmon and steelhead stocks with hatchery fish: A synthesis of published literature. U.S. Dept. Energy, Bonneville Power Administration, Tech. Rep. 90-1. (Available from U.S. Dept. Energy, Bonneville Power Administration, Fish and Wildlife, P.O. Box 3621, Portland, OR 97208.)
- Stillwater Sciences Inc. 1998. North Umpqua Cooperative Watershed Analysis: Synthesis Report Volume 2 [Online]. PacifiCorp, 1998. Available at <http://www.pacificorp.com/File/File986.pdf>. (Accessed January 9 2004.)
- Streamnet. 2002. PNW Reach File. [Vector digital data online]. Available at <ftp://ftp.streamnet.org/pnwr/e00/zip/oreghyd.exe>. (Accessed 7 July, 2003.)

- Sumner, F. H. 1953. Migrations of salmonids in Sand Creek, Oregon. *Trans. Am. Fish. Soc.* 82:139-149.
- Terrel, T. T. 1979. Physical regionalization of coastal ecosystems of the United States and its territories. U.S. Fish and Wildlife Service, Biological Services Program. FWS/OBS-78/80. 30 p.
- U. S. Census Bureau. 2000. 2000 State and State Equivalent Areas Cartographic Boundary Files. [Vector digital data online]. Available at http://www.census.gov/geo/cob/bdy/st/st00shp/st99_d00_shp.zip. (Accessed 25 July, 2003.)
- USGS (U.S. Geological Survey). 2003a. Geographic Names Information System. [Tabular digital data online]. Available at http://geonames.usgs.gov/geonames/stategaz/OR_deci. (Accessed 24 July, 2003.)
- USGS (U.S. Geological Survey). 2003b. Streams and Waterbodies of the United States. [Vector digital data online]. Available at <http://edcftp.cr.usgs.gov/pub/data/nationalatlas/hydrogm020.tar.gz>. (Accessed 18 August 2003.)
- USGS (U.S. Geological Survey). 2003c. Surface Water data for Oregon: Calendar Year Statistics. [Text search online]. Available at <http://nwis.waterdata.usgs.gov/or/nwis/annual>. (Accessed 6 November, 2003.)
- USGS (U.S. Geological Survey). 2003d. Geographic Names Information System. [Tabular digital data online]. Available at http://geonames.usgs.gov/geonames/stategaz/CA_deci. (Accessed 5 Nov., 2003.)
- USGS (U.S. Geological Survey). 2003e. Geographic Names Information System. [Tabular digital data online]. Available at http://geonames.usgs.gov/geonames/stategaz/WA_deci. (Accessed 5 Nov., 2003.)
- Uremovich, B. 1977. Straying of fall Chinook salmon from Elk River Hatchery into Sixes River, 1970-1976. Oregon Dept. of Fish and Wildlife, Research Section, Corvallis, OR.
- Ursitti, V. L. 1989. Residualism of juvenile coho salmon in Eel Lake, Oregon. *Oreg. Dep. Fish Wildl. Fish Div. Info. Rep.* 89-9, 15 p.
- Van Horne, B., and J. A. Wiens. 1991. Forest bird habitat suitability models and development of general habitat models. *U.S. Fish Wildl. Serv., Fish Wildl. Res.* (8). 31 p.
- Waples, R. S. 1990. Conservation genetics of salmon. Vol. II. Effective population size and rate of loss of genetic diversity. *J. Heredity* 81:267-276.
- Waples, R. S. 1991a. Definition of "species" under the Endangered Species Act: Application to Pacific salmon. *U.S. Dep. Commer., NOAA Tech. Memo., NMFS-F/NWC-* 194, 29 p.
- Waples, R. S. 1991b. Pacific salmon, *Oncorhynchus* spp., and the definition of "species" under the Endangered Species Act. *U.S. Natl. Mar. Fish. Serv. Mar. Fish. Rev.* 53(3):11-22.
- Waples, R. S. 1995. Evolutionarily Significant Units and the conservation of biological diversity under the Endangered Species Act. *Am. Fish. Soc. Symp.* 17:8-27.

- Waples, R. S., and D. J. Teel. 1990. Conservation genetics of Pacific salmon. Vol. I. Temporal changes in allele frequency. *Conserv. Biol.* 4:144-156.
- WDF (Washington Department of Fisheries), WDW (Washington Department of Wildlife), and WWTIT (Western Washington Treaty Indian Tribes). 1993. 1992 Washington State salmon and steelhead stock inventory (SASSI). Wash. Dep. Fish Wildl., Olympia, 212 p. plus three appendices: Appendix One–Hood Canal and Strait of Juan de Fuca (December 1994, 424 p.), North Puget Sound (June 1994, 418 p.), and South Puget Sound (September 1994, 371 p.) volumes; Appendix Two–Coastal stocks (August 1994, 587 p.); and Appendix Three–Columbia River stocks (June 1993, 580 p.). (Available Washington Department of Fish and Wildlife, 600 Capitol Way N., Olympia, WA 98501-1091.)
- WDNR (Washington Department of Natural Resources). 2003. Willapa Hills. Available at <http://www.dnr.wa.gov/geology/willapa.htm>. (Accessed date.)
- Weeks, H. (Oregon Department of Fish and Wildlife, Portland, OR). 1994. Letter to L. Weitkamp (NMFS) 5/4/94, Information on Oregon coho salmon life history characteristics.
- Weidemann, A. M. 1984. The Ecology of Pacific Northwest coastal sand dunes: a community profile. U.S. Fish and Wildl. Serv. FWS/OBS-84/04. 130 p.
- Weitkamp, L. A., T. C. Wainwright, G. J. Bryant, G. B. Milner, D. J. Teel, R. G. Kope, and R. S. Waples. 1995. Status review of coho salmon from Washington, Oregon, and California. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-NWFSC-24, 258 p.
- Weitkamp, L., and K. Neely. 2002. Coho salmon (*Oncorhynchus kisutch*) ocean migration patterns: insight from marine coded-wire tag recoveries. *Can. J. Fish. Aquat. Sci.* 59:1100-1115. 16 p.
- Welch, D. W., J. F. T. Morris, M. E. Thiess, M. Trudel, A. R. Ladoucer, M.C. Jacobs, T. B. Zubkowski, and H. R. MacLean. 2004. CCGS W.E. Ricker Gulf of Alaska salmon survey, October 8-27, 2003. Canadian Data Report of Fisheries and Aquatic Sciences 1145.
- Zhou, S. 2000. Stock assessment and optimal escapement of coho salmon in three Oregon coastal lakes. ODFW Fish Division. (Available from Oregon Department of Fish and Wildlife, 3406 Cherry Ave. NE, Salem, OR 97303.)

GLOSSARY

A

Abundance. The number of fish in a *population*.

Active channel width. The distance across a stream or channel as measured from bank to bank at bankfull flow.

Allele. Any one of a number of alternative forms of a gene that can occur at the same location (*locus*) on a chromosome. A *population* can have many alleles for a particular locus, but an individual can carry no more than two alleles at a *diploid* locus.

Allozymes. Alternative forms of an enzyme that have the same function, are produced by different *alleles*, and are often detected by *protein electrophoresis*.

Anthropogenic factor. A circumstance or influence caused or produced by human action.

Artificial propagation. *Hatchery* spawning and rearing of salmon, usually to the *smolt* stage.

AUC (Area Under the Curve). A statistical technique for estimating an annual total number of *spawners* from periodic spawner counts.

B

Barrier. A blockage such as a waterfall, culvert, or rapid that impedes the movement of fish in a stream system.

Biological Review Team (BRT). The team of scientists who evaluate scientific information for the National Marine Fisheries Service (NMFS) status reviews.

C

Carrying capacity. The number of individuals that the resources of a habitat can support.

Catastrophic events. Sudden events that disastrously alter large areas of landscape. These can include floods, landslides, forest fires, and volcanic eruptions.

Channel gradient. The slope of a stream reach.

CIMRS (Cooperative Institute for Marine Resources Studies). A cooperative project between Oregon State University and NOAA Fisheries.

CLAMS (Coastal Landscape Analysis and Modeling Study). A cooperative project between the Oregon State University Department of Forestry and the U.S. Forest Service Pacific Northwest Forest Science Laboratory.

Co-managers. Federal, state, and tribal agencies that cooperatively manage salmon in the Pacific Northwest.

Coded-wire tag (CWT). A small piece (0.25 x 0.5 or 1.0 mm) of stainless steel wire that is injected into the snouts of juvenile salmon and steelhead. Each tag is etched with a binary code that identifies its release group.

D

De-listing. Taking a species off of the endangered species list.

Demographic stochasticity. “Chance events in the survival and reproductive success of a finite number of individuals” (Shaffer 1981).

DEM (Digital Elevation Model). A digital data set representing a topographic map that can be used for computer analysis. We used DEMs to calculate the *intrinsic potential* of stream systems.

Density effects. Survival of juvenile salmon may be influenced by their density. Survival is usually higher when density is low.

Dependent populations. Populations that rely upon immigration from surrounding populations to persist. Without these inputs, dependent populations would have a lower likelihood of *persisting* over 100 years.

Depensation. The effect where a decrease in spawning stock leads to reduced survival or production of eggs through either (1) increased predation per egg given constant predator pressure, or (2) the “Allee effect” (the positive relationship between population density and the reproduction and survival of individuals) with reduced likelihood of finding a mate.

Diploid. Having two complete chromosome pairs in a nucleus.

Distinct population segment (DPS). A *population*, or group of populations of a vertebrate species that is “discrete” from other populations and *significant* to the biological species as a whole.

DNA (deoxyribonucleic acid). A complex molecule that carries an organism’s heritable information. The two types of DNA commonly used to examine genetic variation are *mitochondrial DNA* (mtDNA), a circular molecule that is maternally inherited, and *nuclear DNA*, which is organized into a set of chromosomes (see also *allele* and *electrophoresis*).

Donor populations. These are *Functionally Independent* or *Potentially Independent populations* that are large enough to “donate” *migrant* adults to *Dependent populations*.

E

Ecoregion. An integration of physical and biological factors such as geologic history, climate, and vegetation.

Effective migration rate. The proportion of successfully spawning adults that migrate to a new *population*. (Excludes *migrants* that do not successfully reproduce.) (See also *Migration rate*.)

Electrophoresis. The movement of charged particles in an electric field. This process has been developed as an analytical tool to detect genetic variation revealed by charge differences on proteins or molecular weight in DNA.

Endangered species. A species in danger of extinction throughout all or a significant portion of its range.

ESA. U.S. Endangered Species Act.

Escapement. Usually refers to adult fish that “escape” from both fisheries and natural *mortality* to reach the spawning grounds.

Estuarine habitat. Areas available for feeding, rearing, and smolting in tidally influenced lower reaches of rivers. These include marshes, sloughs and other backwater areas, tidal swamps, and tide channels.

Evaporation potential. The maximum depth of water that could evaporate in a year. Evaporation potential is principally determined by temperature and relative humidity. It is a measure of how much moisture “stress” plants are under due to dry conditions.

Evolutionarily Significant Unit (ESU). An ESU represents a *distinct population segment* of Pacific salmon under the *Endangered Species Act* that 1) is substantially reproductively isolated from conspecific populations and 2) represents an important component of the evolutionary legacy of the species. See also *Distinct population segment*.

Exploitation rate. The proportion of adult fish from a *population* that die as a result of fisheries.

Extinction. The loss of a species, or ESU. May also be used for the extirpation of local populations.

F

Factors for decline. These are factors identified that caused a species to decrease in *abundance* and *distribution* and become threatened or endangered.

Fecundity. The number of offspring produced per female.

Fourth-Field and Fifth-Field Hydrologic Units. In the United States Geological Survey (USGS), hydrologic units have been divided at different scales. The area of a fourth-field hydrologic unit is 440,000 acres and a fifth-field hydrologic unit is between 40,000 and 250,000 acres.

Freshwater habitat. Areas available for spawning, feeding, and rearing in freshwater.

Fry. Young salmon that have emerged from the gravel and no longer have an egg sack.

Functionally Independent population. A high-*persistence population* whose dynamics or extinction risk over a 100-year time frame is not substantially altered by exchanges of individuals with other populations (*migration*). Functionally independent populations are net “donor” populations that may provide *migrants* for other types of populations. This category is analogous to the “independent populations” of McElhaney et al. (2000).

G

Genetic distance. A quantitative measure of the genetic difference between a pair of samples, based on *allele* frequencies at multiple loci.

Genetic drift. Random changes in gene frequencies of *populations*.

Gradient. The slope of a stream system.

H

Habitat quality. The suitability of physical and biological features of an aquatic system to support salmon in the freshwater and estuarine system.

Hatchery. A facility where *artificial propagation* of fish takes place.

Historical abundance. The number of fish that were produced before the influence of European settlement.

Homing fidelity. The propensity of an adult salmon to return to its natal stream.

Hydrology. The distribution and flow of water in an aquatic system.

I

Independent Multidisciplinary Science Team (IMST). A scientific advisory body to the Oregon Legislature and Governor on Watershed, Forestry, Agriculture, and Fisheries Science issues.

Inbreeding depression. Reduced survival rates of individuals in a *population* suffering from the effects of harmful recessive genes through matings between close relatives. Inbreeding depression may become a problem when populations get very small.

Independence. Reflects the interaction between *isolation* and *persistence*. A persistent population that is highly isolated is highly independent.

Intrinsic potential. A modeled attribute of streams that includes the *channel gradient*, *valley constraint* and *mean annual discharge of water*. Intrinsic potential in this report refers to a measure of potential coho salmon habitat quality (Burnett et al. 2003).

Isolating mechanisms. Things that reduce the ability of *populations* to interbreed. These could include physical mechanisms such as distance, and behavioral mechanisms such as *run timing*.

Isolation. The degree to which a *population* is unaffected by *migration* to and from other populations. As the influence of migration decreases, a population's isolation increases.

J

Jack. A coho salmon that matures at age 2 and returns from the ocean to spawn a year earlier than normal. Jacks are all male fish.

Jacking rate. The proportion of adult coho salmon from a brood that return as *jacks*.

Juvenile. A fish that has not matured sexually.

L

Latitudinal cline (also known as a latitudinal diversity gradient.) A change in a biological trait that occurs across latitudes. For example, *jacking rate* generally increases from north to south.

Life history. The specific life cycle of a fish from egg to adult.

Limiting factors. Factors that limit survival or *abundance*. They are usually related to habitat quantity or quality at different stages of the life cycle. Harvest and predation may also be limiting factors.

Listed species. Species included on the “List of Endangered and Threatened Species” authorized under the *Endangered Species Act* and maintained by the *U.S. Fish and Wildlife Service* and *NOAA Fisheries*.

Littoral zone. In lakes, the area of lake bottom that receives enough light for rooted plants to grow. In the ocean, the marine ecological realm that experiences the effects of tidal and longshore currents and breaking waves to a depth of 5 to 10 m (16 to 33 feet) below the low-tide level, depending on the intensity of storm waves. (Encyclopædia Britannica 2004).

Locus. Location on a chromosome that holds a specific gene. Plural is loci.

Lowland habitat. Low-gradient stream habitat with slow currents, pools, and backwaters used by fish. This habitat is often converted to agricultural or urban use.

M

Marine survival rate. The proportion of smolts entering the ocean that return as adults.

Mean annual discharge of water. A single value or average that summarizes or represents the annual discharge amount, typically expressed in cubic meters per second.

MES, Inc. Manufacturing Engineering Systems, Inc.

Metric. A unit of measure.

Microsatellite. A class of repetitive DNA used for estimating genetic distances.

Migrant. A fish that is born in one *population* but returns to another population to spawn.

Migration. Movement of fish from one *population* to another.

Migration rate. The proportion of *spawners* that migrate from one population to another. See also *Effective migration rate*.

Mitochondrial DNA (mtDNA). The *DNA* genome contained within mitochondria and encoding a small subset of mitochondrial functions. Only female mtDNA is transmitted to the next generation.

Mortality. Death.

N

NMFS. National Marine Fisheries Service, also known as NOAA Fisheries.

NOAA. National Oceanic and Atmosphere Administration.

NOAA Fisheries. The fisheries branch of NOAA, also known as NMFS.

Nuclear DNA (nDNA). The DNA contained in the chromosomes within the nucleus of eukaryotic cells. The nuclear genome in Pacific salmon is approximately 4.6 billion base pairs in size.

NWFSC. NOAA Fisheries Northwest Fisheries Science Center.

O

ODFW. Oregon Department of Fish and Wildlife.

ONCC. Oregon Northern California Coho Salmon Technical Recovery Team.

ORISE. Oak Ridge Institute for Science and Education.

OWEB. Oregon Watershed Enhancement Board.

P

Parr. The life stage of salmonids that occurs after *fry* and is generally recognizable by dark vertical bars (parr marks) on the sides of the fish.

Persistence. A *population's* relative ability to sustain itself without input from neighboring populations.

Phylogenetic tree. The genetic distances and relationships among *populations*, frequently represented with a branching or tree-like diagram.

Population. A group of fish of the same species that spawns in a particular locality at a particular season and does not interbreed substantially with fish from any other group.

Population classification. The grouping of *populations* into *Functionally Independent*, *Potentially Independent*, and *Dependent* classes.

Population dynamics. Changes in the number, age, and sex of individuals in a *population* over time, and the factors that influence those changes. Five components of populations that are the basis of population dynamics are birth, death, sex ratio, age structure, and dispersal.

Population identification. Delineating the boundaries of *historical populations*.

Population structure. This includes measures of age, density, and growth of fish populations.

Potentially Independent populations. *High-persistence populations* whose *population dynamics* are substantially influenced by periodic immigration from other populations. In the event of the

decline or disappearance of *migrants* from other populations, a Potentially Independent population could become a *Functionally Independent* population.

Production. The number of fish produced by a *population* in a year.

Productive capacity. Maximum possible *production* from a given area.

Productivity. The rate at which a *population* is able to produce fish.

Protein electrophoresis. An analytical laboratory technique that measures differences in the amino acid composition of proteins from different individuals. See also *Electrophoresis*.

R

Recovery. The reestablishment of a threatened or endangered species to a self-sustaining level in its natural ecosystem (in other words, to the point where the protective measures of the Endangered Species Act are no longer necessary).

Recovery domain. The area and species that the Technical Recovery Team is responsible for.

Recovery plan. A document identifying actions needed to make *populations* of naturally produced fish comprising the Oregon Coast Coho Salmon *ESU* sufficiently *abundant*, *productive*, and diverse so that the *ESU* as a whole will be self-sustaining and will provide environmental, cultural, and economic benefits. A recovery plan will also include goals and criteria by which to measure the *ESU*'s achievement of recovery, and an estimate of the time and cost required to carry out the actions needed to achieve the plan's goals.

Recovery scenarios. Various sequences of events expected to lead to *recovery* of Oregon Coast Coho Salmon.

Run timing. The time of year (usually identified by week) when spawning salmon return to the spawning beds.

S

Salmonids. Any of the species included in salmon, trout, and char.

Significant. Biological significance refers to an effect that has a noteworthy impact on health or survival.

Smolt. A life stage of salmon that occurs just before the fish leaves freshwater. Smolting is the physiological process that allows salmon to make the transition from fresh to salt water.

Smolt capacity. The maximum number of smolts a basin can produce. Smolt capacity is related to habitat quantity and quality.

Spawners. Adult fish on the spawning grounds.

Species. Biological definition: A small group of organisms formally recognized by the scientific community as distinct from other groups. Legal definition. Refers to joint policy of the USFWS and NMFS that considers a species as defined by the ESA to include biological species, subspecies, and *DPSs*.

Stock transfer. The practice of moving fish between basins or *populations*.

Stray rate. As used in this document, the stray rate refers to the number of spawning adults that return to a stream other than their natal stream within a basin. (See also *Migration rate*.)

Sustainability. The ability of a stock to *persist* or sustain itself over long periods of time.

SWFSC. NOAA Southwest Fisheries Science Center.

T

Threatened species. A species not presently in danger of extinction but likely to become so in the foreseeable future.

TRT. Technical Recovery Team.

U

USFS. United States Forest Service.

USGS. United States Geologic Survey.

V

Valley constraint. The valley width available for a stream or river to move between valley slopes.

Viability. The likelihood that a *population* will *sustain* itself over a 100-year time frame.

Viability criteria. A prescription of a *population* conservation program that will lead to the *ESU* having a negligible risk of extinction over a 100-year time frame.

W

Warmwater fish. Spiny-rayed fish such as sculpins, minnows, darters, bass, walleye, crappie, and bluegill that generally tolerate or thrive in warm water.